

Model for plasma jet-driven magneto-inertial fusion

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2017 J. Phys.: Conf. Ser. 875 122002

(<http://iopscience.iop.org/1742-6596/875/13/122002>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 195.19.39.67

This content was downloaded on 30/08/2017 at 11:28

Please note that [terms and conditions apply](#).

You may also be interested in:

[Status of the US program in magneto-inertial fusion](#)

Y C F Thio

[Finite- Effect on the Dynamics of Double Gas-Puff Staged Pinch](#)

Farah Deeba, Kamaluddin Ahmed, Mahnaz Qader Haseeb et al.

[50 years of fusion research and the road to fusion energy](#)

M. Kikuchi

[50 years of fusion research and the road to fusion energy](#)

Werner Burkart

[Optimization of Shell Implosion Driven by Black Body Radiation](#)

Masakatsu Murakami and Katsunobu Nishihara

[Comment on "On quantum plasma: A plea for a common sense" by Vranjes J. et al.](#)

P. K. Shukla and M. Akbari-Moghanjoughi

[Joint Varenna-Lausanne International Workshop on the Theory of Fusion Plasmas 2016](#)

[START plasma overcomes large-scale instabilities](#)

Jim Drake and Adil Hassam

[Thermonuclear fusion with a spinning gas-puff staged pinch](#)

Arshad M Mirza, N A D Khattak, M Salahuddin et al.

Model for plasma jet-driven magneto-inertial fusion

Sergei V. Ryzhkov*¹, Victor V. Kuzenov*^{†‡} ²

* Department of Thermal Physics, Bauman Moscow State Technical University, Moscow 105005, Russia

[†] N.L. Dukhov All-Russian Research Institute of Automatics (State Corporation «ROSATOM»), Moscow 127055

[‡] A.Yu. Ishlinsky Institute for Problems in Mechanics RAS, Moscow 119526, Russia

Synopsis Mathematical models are developed to model the implosion dynamics for plasma jets-magnetized target system [1-9]. Transport (diffusion, thermal, radiative) processes of atoms, ions, electrons, molecules, photons in a strong magnetic field are taken into account.

We assume that the plasma is quasi-neutral, and use the average ion model to determine Z_i (when it is assumed that all ions of chemical components "i" have the same value Z_i at each point in the computational domain). This can be justified if the distribution of the different degree of ionization is sufficiently narrow.

The average ion charge $Z_i(t)$ can change over time and we can use the formula:

$$\frac{DZ_i}{Dt} = Z_i(v_I - v_{Ph} - v_{T.R}).$$

The following inelastic processes (electron impact ionization v_I , photorecombination v_{Ph} and triple recombination $v_{T.R}$) in the one-component case are taken into consideration in calculation:

$$v_I = 10^{-7} n_i \left[\frac{I_H}{I(Z_i)} \right]^{3/2} \exp \left[-\frac{I(Z_i)}{kT_e} \right] \left[\frac{I(Z_i)}{kT_e} \right]^{1/2} \left[1 + \frac{I(Z_i)}{kT_e} \right]^{-1}$$

$$v_{Ph} = 6 \times \lambda \times 10^{-14} n_i \left[\frac{I(Z_i)}{I_H} \right]^{1/2} \left[\frac{I(Z_i)}{kT_e} \right]^{3/2} \left[1 + \frac{I(Z_i)}{kT_e} \right]^{-1}$$

$$v_{T.R} = 3.3 \times 10^{-31} Z_i n_i^2 \left[\frac{I_H}{I(Z_i)} \right]^{3/2} \left[\frac{I_H}{kT_e} \right]^{3/2} \left[\frac{I(Z_i)}{kT_e} \right]^{1/2} \left[1 + \frac{I(Z_i)}{kT_e} \right]^{-1}$$

where $I_H = 13.6$ eV is the ionization potential of hydrogen, $I(Z_i)$ is the ionization energy of the average ion, T_e is the electron temperature (K), $\lambda = 1 \div 5$, $n_e = Z_i n_i$, $n_i = \rho/M_i$ are electron and ion densities (cm^{-3}).

It is known, that plasma dynamical processes are strongly affected by irradiation, if plasma temperature achieve 1 eV. In that case, the gas dynamic fields of thermal physical variables may be calculated only with taking into account the radiation fields. In this work the radiation transfer equation is used in the form of multi-group diffusion approach:

$$\frac{1}{J} \frac{\partial (J q_{i\xi})}{\partial \xi} + \frac{1}{J} \frac{\partial (J q_{i\eta})}{\partial \eta} + \chi_i c U_i = 4 \chi_i \sigma_i T^4,$$

$$\frac{c}{3} \frac{\partial U_i}{\partial \xi} + \chi_i q_{i\xi} = 0, \quad \frac{c}{3} \frac{\partial U_i}{\partial \eta} + \chi_i q_{i\eta} = 0,$$

where $U_i(y, z, t)$ is the radiation power density in i -th spectral group, χ_i is the spectral absorption coefficient. Besides of the method mentioned above, the discrete ordinates method has been used in the work, which gets an opportunity to solve the radiation transfer equation on tetrahedral non-structured mesh. Modified alternatively triangular three-layered iterative scheme is applied for the solution of radiation transport equations, where the time step selected via conjugate directions method.

The turbulent viscous coefficient $\mu_{\Sigma}^{r,z}$ is calculated by using of the Bussinesk hypothesis, when effective viscous is $\mu_{\Sigma}^{r,z} = \mu_m^{r,z} + \mu_t$, where $\mu_m^{r,z}$ is a molecular viscosity coefficient, taking into account the atomic-molecular collision processes and the presence of a magnetic field in the plasma, and μ_t is a turbulent one, determined from two-parameter turbulence model (Coakley model). The mathematical model and test results are presented and discussed.

This research has been supported by the Ministry of Education and Science of the Russian Federation (Project No. 13.5240.2017/БЧ).

References

- [1] H. Nagatomo *et al.* 2016 *J. Phys. Conf. Ser.* **717** 012041
- [2] V.V. Kuzenov *et al.* 2013 *Probl. At. Sci. Technol.* **4** (86) 103
- [3] S.V. Ryzhkov *et al.* 2013 *Fusion Sci. Technol.* **63** (1T) 135
- [4] V.Kuzenov *et al.* 2013 *Probl. At. Sci. Tech.* **1** 12
- [5] S. Ryzhkov 2010 *Probl. At. Sci. Technol.* **4** 105
- [6] S.A. Belkov *et al.* 1997 *JETP* **84** 272
- [7] V.V. Kuzenov *et al.* 2016 *Bull. Russ. Acad. Sci.: Phys.* **80** 598
- [8] I.Yu. Kostyukov *et al.* 2011 *Plasma Phys. Rep.* **37** 1092
- [9] V.V. Kuzenov *et al.* 2017 *J. Phys. Conf. Ser.* **815** 012024

¹ E-mail: vik.kuzenov@gmail.com ^P

² E-mail: svryzhkov@bmstu.ru

